P38 Polyclonal Antibody

Catalog No. E-AB-32459

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse,Rat
Immunogen	Synthesized peptide derived from human p38 around the non-phosphorylation site of Thr180/Tyr182.
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide, 0.5% BSA and 50% glycerol, pH7.4

Applications — Recommended Dilution

WB 1:500-1:2000,
IHC 1:100-1:300,
ELISA 1:5000

Data



Western Blot analysis of HepG2, 293-UV cells using p38 Polyclonal Antibody at dilution of 1:3000.
Observed Mw:38kDa
Calculated Mw:41kDa

Preparation & Storage

Storage	Store at -20℃. Avoid freeze / thaw cycles.

Background

Responds to activation by environmental stress, pro-inflammatory cytokines and lipopolysaccharide (LPS) by phosphorylating a number of transcription factors, such as ELK1 and ATF2 and several downstream kinases, such as MAPKAPK2 and MAPKAPK5. Plays a critical role in the production of some cytokines, for example IL-6. May play a role in stabilization of EPO mRNA during hypoxic stress. Isoform Mxi2 activation is stimulated by mitogens and oxidative stress and only poorly phosphorylates ELK1 and ATF2. Isoform Exip may play a role in the early onset of apoptosis.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com