PKM Polyclonal Antibody

Catalog No. E-AB-13771


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human,Mouse,Rat
Immunogen	Recombinant protein of human PKM2
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Applications	Recommended Dilution
WB	1:500-1:2000
IHC	1:50-1:200

Data



Western Blot analysis of Hela cell using PKM Polyclonal Antibody at dilution of 1:450
Calculated Mw:58kDa



Immunohistochemistry of paraffin-embedded Human cervical cancer using PKM Polyclonal Antibody at dilution of 1:50

Preparation & Storage

Storage Store at -20℃. Avoid freeze / thaw cycles.

Background

This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP, generating ATP and Stimulates POU5F1-mediated transcriptional activation. Plays a general role in caspase independent cell death of tumor cells. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial pathogenesis.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com