

NUP62 Polyclonal Antibody

Catalog No. E-AB-15263

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Reactivity	Human,Mouse,Rat
Immunogen	Recombinant protein of human NUP62
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Applications Recommended Dilution


WB	1:500-1:2000
IHC	1:50-1:200

Data



Western Blot analysis of Jurkat cell and Human brain malignant glioma tissue, PC3, K562 and A549 cell using NUP62 Polyclonal Antibody at dilution of 1:300

Calculated Mw:53kDa



Immunohistochemistry of paraffin-embedded Human liver cancer using NUP62 Polyclonal Antibody at dilution of 1:40

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

The nuclear pore complex is a massive structure that extends across the nuclear envelope, forming a gateway that regulates the flow of macromolecules between the nucleus and the cytoplasm. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells. The protein encoded by this gene is a member of the FG-repeat containing nucleoporins and is localized to the nuclear pore central plug. This protein associates with the importin alpha/beta complex which is involved in the import of proteins containing nuclear localization signals. Multiple transcript variants of this gene encode a single protein isoform.

For Research Use Only