PRKD3 Polyclonal Antibody

Catalog No. E-AB-15940


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human,Mouse
Immunogen	Synthetic peptide of human PRKD3
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Buffer	PBS with 0.05% sodium azide and 50% glycerol, PH7.4

Applications	Recommended Dilution
WB	1:200-1:1000
IHC	1:50-1:200


Data



Western Blot analysis of Human testis tissue, K562, A549, Raji, NIH/3T3 and Hela cell using PRKD3 Polyclonal Antibody at dilution of 1:200
Calculated Mw:100kDa



Immunohistochemistry of paraffin-embedded Rat liver using PRKD3 Polyclonal Antibody at dilution of 1:200



Immunohistochemistry of paraffin-embedded Human colon cancer using PRKD3 Polyclonal Antibody at dilution of 1:50

Preparation & Storage

Storage	Store at -20℃. Avoid freeze / thaw cycles.

Toll-free: 1-888-852-8623
Tel: 1-832-243-6086
Fax: 1-832-243-6017
Web: www.elabscience.com
Email: techsupport@elabscience.com

Background

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. EacHuman, Mouseember of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family members.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com