PAIP1 Polyclonal Antibody

Catalog No. E-AB-40457


Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description


Reactivity	Human,Mouse,Rat
Immunogen	Recombinant Human Polyadenylate-binding protein-interacting protein 1 protein
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Conjugation	Unconjugated
Buffer	PBS with 0.02% sodium azide,1% protective protein and 50% glycerol,pH7.4

Applications	Recommended Dilution
IHC	1:100-1:300


Data




Immunohistochemistry of paraffin-embedded Human testis using PAIP1 Polycloanl Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Human kidney using PAIP1 Polycloanl Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Mouse kidney using PAIP1 Polycloanl Antibody at dilution of 1:200.



Immunohistochemistry of paraffin-embedded Rat kidney using PAIP1 Polycloanl Antibody at dilution of 1:200.

Preparation & Storage

Storage Store at -20℃. Avoid freeze / thaw cycles.

Background

The protein encoded by this gene interacts with poly(A)-binding protein and with the cap-binding complex eIF4A. It is

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com

involved in translational initiation and protein biosynthesis.Among its related pathways are Deadenylation-dependent mRNA decay and Gene Expression. Overexpression of this gene in COS7 cells stimulates translation. Alternative splicing occurs at this locus and three transcript variants encoding three distinct isoforms have been identified. Gene Ontology (GO) annotations related to this gene include RNA binding and translation activator activity. An important paralog of this gene is CTIF.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com